


Supplementary Figures

Table S1: Characteristics of people diagnosed with OG cancer by tumour type, 2020-2022

	Oes SCC	Oes ACA	Stomach
Male (%)	48.0%	80.2%	66.3%
Median age (yrs)	72	72	74
Age group			
<60	14.7%	16.2%	17.7%
60-69	26.4%	25.4%	19.4%
70-79	34.8%	36.6%	32.3%
≥80	24.1%	21.8%	30.6%
Clinical stage (pre-treatment)			
Stage 0/1	2.7%	4.4%	10.8%
Stage 2	26.5%	8.3%	17.4%
Stage 3	38.7%	40.1%	26.6%
Stage 4	32.1%	47.2%	45.2%
Missing	533	1,515	892
Deprivation quintile (IMD)			
1 – least deprived	17.1%	19.6%	17.3%
2	19.6%	21.7%	18.7%
3	21.2%	21.8%	20.2%
4	20.4%	18.8%	21.7%
5 – most deprived	21.7%	18.1%	22.1%
Missing	28	42	19
Any comorbidities	60.5%	59.1%	60.3%
Performance status			
0 – normal activities	33.9%	40.2%	33.3%
1	36.2%	33.6%	34.7%
2	18.4%	15.6%	18.1%
3	9.9%	9.1%	11.7%
4 – completely disabled	1.6%	1.5%	2.1%
Total	3,646	11,192	5,027

KEY: Oes – oesophageal; SCC – squamous cell carcinoma; ACA – adenocarcinoma; IMD – Index of Multiple Deprivation

Figure S1: Adjusted rates of patients diagnosed after an emergency admission by Cancer Alliance / Welsh region. Graph shows adjusted rates with 95% confidence interval (CI)



NOTE: Rates adjusted for age, sex, deprivation, site of cancer and presence of comorbidities