


Report at a glance

COPD

During 2023–25

Patients diagnosed with COPD in the last 2 years who have a post-bronchodilator spirometry code available in the last 2 years

23.5%



Patients with COPD who are breathless (MRC score 3–5) and have been referred to pulmonary rehabilitation in the last 3 years

16.4%

Patients with COPD who **did not have a record of their smoking status** in the last 15 months

29.6%



Patients with COPD who **did not have a record of their vaping status** in the last 15 months

98.1%

During 2023–25


Adults diagnosed with asthma who have a record of at least one objective measurement* in the last 2 years

62.1%



Adults with asthma who had a personalised asthma action plan (PAAP) in the last 15 months

28.7%



Adults with asthma **who did not have a record of their smoking status** in the last 15 months

35.5%

Adults with asthma **who did not have a record of their vaping status** in the last 15 months

98.2%



CYP asthma

During 2023–25

Children and young people (CYP) diagnosed with asthma who have a record of at least one objective measurement* in the last 2 years

47.4%



CYP (aged 14–18 years) with asthma **who did not have a record of their smoking status** in the last 15 months

65.4%



CYP with asthma who had a personalised asthma action plan (PAAP) in the last 15 months

24.0%



CYP (aged 14–18 years) with asthma **who did not have a record of their vaping status** in the last 15 months

98.6%

*One objective measurement includes spirometry, peak flow with greater than one reading, or evidence of peak flow diary or recorded FeNO